

Prevalence, Associated Factors, and Help-Seeking Behavior of Psychological Distress among International Students at the National University of Malaysia

Khadiga Kahwa, Aniza Ismail

Abstract—Depression, anxiety, and stress are associated with decreased role functioning, productivity, and quality of life. International students are more prone to psychological distress as they face many stressors while studying abroad. The objectives of the study were to determine the prevalence and associated factors of depression, anxiety, and stress among international students, their help-seeking behavior, and their awareness of the available on-campus mental support services. A cross-sectional study with a purposive sampling method was performed on 280 international students at Universiti Kebangsaan Malaysia (UKM) between the age of 18 and 35 years. The Depression Anxiety Stress Scale-21 (DASS-21) questionnaire was used anonymously to assess the mental health of students. Socio-demographic, help-seeking behavior, and awareness data were obtained. Independent sample t-test, one-way ANOVA test, and multiple linear regression were used to explore associated factors. The overall prevalence of depression, anxiety, and stress among international students were 58.9%, 71.8%, and 53.9%, respectively. Age was significantly associated with depression and anxiety. Ethnicity showed a significant association with depression and stress. No other factors were found to be significantly associated with psychological distress. Only 9.6% of the international students had sought help from on-campus mental support services. Students who were aware of the presence of such services were only 21.4% of the participants. In conclusion, this study addressed the gap in the literature on the mental health of international students and provided data that could be used in intervention programs to improve the mental health of the increasing number of international students in Malaysia.

Keywords—Anxiety, depression, stress, help-seeking behavior, students.

I. INTRODUCTION

DEPRESSION, anxiety, and stress are extremely prevalent among university students and their prevalence appears to be increasing [1]-[3]. These psychological problems have a negative impact on the students' quality of life, which in turn can interfere with the healthy adaptation and coping of the international students with the new environment. Psychological distress (PD) is a state of psychological discomfort usually experienced as sadness, depression, stress and anxiety [4], [5]. It often co-exists with somatic

Khadiga Kahwa is Master student in the department of community health, Faculty of medicine, The National University Malaysia, Kuala Lumpur.

Aniza Ismail is with Department of community health, Faculty of medicine National University Malaysia, Kuala Lumpur (e-mail: aniza@ppukm.ukm.edu.my).

complaints, chronic conditions and syndromes that have no medical explanation [6]. Risk factors for PD include stress-related factors and sociodemographic factors [5] such as; age, gender, race, marital status, socioeconomic level and study program [7].

International students are a high-risk group of people who need to be connected with care and support as it is very important to help them develop effective coping strategies for their mental health problems. Since international students come from different cultures and environments, care should be given to increase their awareness about mental illness and the professional help-seeking behavior for PD. It was demonstrated that more than two-thirds of young people did not seek help for their psychological problems [8]-[10]. Sometimes, students may not seek help from professionals since they are unaware of the presence of mental support services at their universities [11]. It has been historically reported that international students underutilized on-campus mental support services [12], [13].

Malaysia is a popular destination for international students worldwide. It was recognized by the United Nations Educational, Science and Cultural Organization (UNESCO) as one of the most 10 preferred destinations for post-secondary education among international students [14]. From a public health perspective, the early detection and prevention of mental health disorders among international university students is very crucial. Although previous studies have been conducted in Malaysia to assess prevalence and associated factors of depression, anxiety and stress among university students [7], [19], more studies are still needed to explore these associated factors, as well as the help-seeking behavior of PD among international university students in Malaysia. This will help to develop and evaluate more effective preventive and therapeutic interventions to these conditions and to encourage students to seek medical help.

II. MATERIALS AND METHODS

A. Study Design

Cross-sectional study was performed on 280 international students at UKM between the ages of 18 and 35 years.

B. Sampling Method

Purposive sampling method was used to select the sample

population of this study.

C. Conflict of interest

The investigators declare that there is no conflict of interest.

D. Methods

Participants were given hardcopies of a self-administered questionnaire which consisted of four sections: DASS-21 questionnaire, socio-demographic data, study program, and lastly, help seeking-behavior and awareness of on-campus mental support services. DASS-21 measures the characteristic and severity of a range of symptoms of depression, anxiety and stress [15]. DASS 21-item, which is a modified version of the original Scale of DASS-42, has many advantages as it is more convenient for people with poor concentration since it is shorter [16]. Cronbach's alpha scores showed reliability rates for the depression, anxiety, and stress scales at 0.91, 0.84, and 0.90 respectively in the normative sample [7]. The questionnaire consists of 21 items, a set of three self-report scales to measure the negative emotional states of depression, anxiety, and stress, as well as seven items for each psychological state [15]. The students were asked to score the extent of various symptoms they experienced over the past week. Each item was scored on a 4-point Likert scale (0 = Did not apply to me at all, 1 = Applied to me to some degree, or some of the time, 2 = Applied to me to a considerable degree, or a good part of time, and 3 = Applied to me very much, or most of the time) [7]. According to the manual guidelines [15], scores from each subscale were summed up and multiplied by two to meet the original 42 items. Each subscale score ranged between 0-42 and the severity scores are presented in Table I. Higher scores indicated greater levels of PD [15].

TABLE I
LOVIBOND SCORING SCALE FOR PSYCHOLOGICAL MOOD

	Stress scale	Anxiety scale	Depression scale
Normal	0-14	0-7	0-9
Mild	15-18	8-9	10-13
Moderate	19-25	10-14	14-20
Severe	26-33	15-19	21-27
Very severe	≥34	≥20	≥28

The sociodemographic data section consisted of five questions to gather information the participants such as age, gender, ethnicity, marital status, and monthly income. The study program section contained one question about study program and educational level of the participants while the help-seeking behavior and awareness of on-campus mental support services section included two dichotomous questions regarding help-seeking behavior for PD and awareness of the presence of on-campus mental support services.

E. Data Analysis

Data analysis was performed using IBM Statistical Package for Social Sciences (SPSS) version 23. Data cleaning was done to remove any missing values. Age, gender, ethnicity, marital status, study program, socioeconomic level, help-seeking behavior and awareness were presented in number and

percentage while the outcome variables such as depression, anxiety, and stress scores were presented as mean and standard deviation. The independent samples t-test and the one-way ANOVA test were used to test the differences in means of depression, anxiety, and stress scores with study variables. Multiple Linear Regression was performed to assess the association of depression, anxiety and stress scores with all factors studied. We consider p value less than 0.05 to be statistically significant, with 95% of confidence interval.

III. RESULTS AND DISCUSSION

A. Characteristics of Study Participants

Table II represents the characteristics of study participants. The total number of participating respondents was 280 students.

TABLE II
CHARACTERISTICS OF THE STUDY PARTICIPANTS

Variables	Study participants (n=280)	
	Number	Percent (%)
Age Group:		
18-20 years	25	8.9
21-24 years	60	21.4
25-35 years	195	69.6
Gender:		
Female	66	23.6
Male	214	76.4
Nationality:		
Africans	20	7.1
Arabs	168	60.0
Asians	83	29.6
Others	9	3.2
Educational level:		
Undergraduate	43	15.4
Postgraduate	237	84.6
Study Program:		
Non-medical	237	84.6
Medical/paramedical	43	15.4
Monthly Income:		
< 2000 RM	140	50.0
2000 – 5000 RM	115	41.1
> 5000 RM	25	8.9
Marital Status:		
Married	112	40.0
Single	168	60.0

B. Prevalence of Depression, Anxiety and Stress among Study Participants

TABLE III
PREVALENCE OF DEPRESSION, ANXIETY, AND STRESS AMONG STUDY PARTICIPANTS

Variables	Depression		Anxiety		Stress	
	Number	%	Number	%	Number	%
Levels:						
Normal	115	41.1	79	28.2	129	46.1
Mild	34	12.1	25	8.9	44	15.7
Moderate	73	26.1	58	20.7	48	17.1
Severe	34	12.1	45	16.1	52	18.6
Very severe	24	8.6	73	26.1	7	2.5
Score:						
Mean ± SD	13.1 ± 9.2		13.8 ± 9.2		16.6 ± 9.0	
Median	12.0		12.0		16.0	
Min – Max.	0.0 – 42.0		0.0 – 42.0		0.0 – 42.0	

Table III reports the prevalence of depression, anxiety, and stress and their associated severity among study participants. Depression, anxiety, and stress were divided according to

severity into five groups which are; normal, mild, moderate, severe and very severe. The overall prevalence of depression, anxiety, and stress among international students were 58.9%, 71.8% and 53.9%, respectively.

C. Factors Associated with Depression, Anxiety and Stress among Study Participants

As shown in Table IV, depression and anxiety scores were significantly associated with age. Higher depression and anxiety scores were found among the age group of 21-24

years. The stress scores were not found to be significantly different by age group. However, the depression and stress scores show significant association with ethnicity, with higher depression and stress scores being recorded in the minority ethnic groups, followed by Arabs, Asians and lastly, Africans. While results show that the factors of gender, study program, monthly income, and marital status were not significantly associated with depression, anxiety, and stress.

TABLE IV
ASSOCIATION BETWEEN DEPRESSION, ANXIETY AND STRESS AND CHARACTERISTICS OF STUDY PARTICIPANTS

Variables	Depression score			Anxiety score			Stress score		
	Mean ± Standard deviation (SD)	Test of significance	P-value P < 0.05	Mean ± Standard deviation (SD)	Test of significance	P-value P < 0.05	Mean ± Standard deviation (SD)	Test of significance	P-value P < 0.05
Age group									
18-20	11.3 ± 6.8	one-way ANOVA	0.042	12.3±6.8	one-way ANOVA	0.014	16.2±7.9	one-way ANOVA	0.303
21-24	15.6 ± 8.5			16.8±6.7			18.2 ± 8		
25-35	12.5 ± 9.5			13 ± 9.5			16.2±9.4		
Gender									
Male	13 ± 9.3	t-test	0.918	13.9±9.4	t-test	0.755	16.3 ± 9	t-test	0.289
Female	13.2 ± 8.7			-0.103			13.5±8.6		
Ethnicity									
Arabs	14.1± 9.5	one-way ANOVA	0.01	14.4 ± 9.6	one-way ANOVA	0.231	17.3±9.5	one-way ANOVA	0.027
Asians	12.1± 8.1			13.5 ± 7.9			16.6±7.4		
Africans	7.3 ± 7.8			10 ± 8.7			10.9±9.1		
Others	14.9±10.6			12.7±13.6			18 ± 11		
Education level									
Undergraduate	12.7 ± 8	t-test	0.785	13.6 ± 8.3	t-test	0.888	16.2±7.4	t-test	0.712
Postgraduate	13.1 ± 9.4			-0.273			13.8 ± 9.4		
Study program									
Non-medical	12.9 ± 9.2	t-test	0.603	13.8 ± 9.4	t-test	0.945	16.5 ± 9	t-test	0.705
Medical/Paramedical	13.7 ± 8.9			-0.521			13.7 ± 8.3		
Monthly income									
< 2000MYR	12.8 ± 8.9	one-way ANOVA	0.924	13.6 ± 8.7	one-way ANOVA	0.787	16.8±8.9	one-way ANOVA	0.824
2000-5000MYR	13.3 ± 9.5			13.7±10.1			16.3±9.4		
> 5000MYR	13 ± 9.3			0.079			14.7 ± 8.4		
Marital status									
Married	12.3 ± 9.8	t-test	0.267	13.6 ± 9.7	t-test	0.837	16.8±9.5	t-test	0.863
Single	13.5 ± 8.8			-1.113			13.9 ± 8.9		

Multiple linear regression was performed and the results are presented in Table VI. Depression was found to be significantly related with age group and ethnicity while each of anxiety and stress were found to be related to only one variable which was the age group and ethnicity, respectively.

D.Help-Seeking Behavior and Awareness of On-Campus Services among Study Participants

Help-seeking behavior among international students was represented in Fig. 1. It shows that only 9.6% of students had sought help for their PD from the available university services. As shown in Table V, depression, anxiety, and stress scores were significantly associated with help-seeking behavior. Students who had higher depression, anxiety and stress scores were more likely to seek professional help while students' awareness of the presence of on-campus mental support services was only 21.4%, as shown in Fig. 2.




Fig. 1 Help seeking for depression, anxiety and stress from UKM health services

E. Discussion

The present study assessed the PD among international university students in Malaysia by determining the prevalence of depression, anxiety, and stress and examining their associated risk factors among this population. In addition, it

evaluated their help-seeking behavior and the awareness about the presence of on-campus mental support services.




Fig. 2 Awareness of available on-campus mental health services

TABLE V
ASSOCIATION BETWEEN DEPRESSION, ANXIETY AND STRESS AND HELP-SEEKING BEHAVIOR

	Help-seeking behavior: (Mean \pm standard deviation)		Test of significance	P-value ($P < 0.05$)
	Yes	No		
Depression score	19.11 \pm 7.4	12.40 \pm 9.1	t-test	0.00
Anxiety score	20.3 \pm 8.97	13.1 \pm 8.98		0.00
Stress score	22.15 \pm 9.36	16.04 \pm 8.80		0.001

The results of the present study showed high prevalence and severity of depression, anxiety, and stress among international students with more than half of this population suffering from mild to very severe levels of PD. Anxiety appeared to be the most prevalent with more than two thirds of the students reporting they had some sort of anxiety disorders. The prevalence of moderate to very severe depression, anxiety, and stress were found to be 46.8%, 62.9% and 38.2%, respectively. This prevalence is higher compared to the prevalence of depression, anxiety, and stress of moderate severity and above recorded among Malaysian and Turkish university students at rates of 37.2%, 63% and 23.7% and 27.1%, 47.1% and 27%, respectively [17]. The overall prevalence of stress in this study was consistent with that of an earlier study among international students at 54% [7]. Higher prevalence of PD among international students may be because they have to adapt to the new environment and different education system, they face linguistic barriers and changes in their support system. Moreover, social stressors such as cultural difference, relationships with peers and hostel mates may affect their mental well-being. These findings are in accordance with the study hypothesis that there is high prevalence and severity of depression, anxiety and stress among international university students in Malaysia. Although DASS-21 is a screening and not a diagnostic tool, this high prevalence of depression, anxiety and stress may need further consideration from university administrative staff and health-care professionals.

In this study, age was found to have significant associations with depression (p-value is 0.042) and anxiety (p-value is 0.014) scores. This is consistent with other studies which reported that age was significantly associated with depression and anxiety scores [7], [18]. Higher depression and anxiety

scores were detected in the age group of 21-24 years followed by 25-35 years, and lastly, younger students aged 18-20 years. This is consistent with a previous study which showed that age more than 20 years has strong association with depression, anxiety and stress [18]. This may be because older students are more worried about their future and life after graduation. Older students may be keener with regard to finishing their studies, so as to be able to work and be financially independent.

This study revealed that there were no significant associations between gender and depression, anxiety, or stress, with both male and female students reporting almost similar levels of PD. This is in accordance with a previous study which showed that gender had no significant association with depression, anxiety, and stress among university students [19]. On the other hand, other studies found that females had higher levels of PD than males [17], [20]. It was demonstrated that females were twice as likely as males to have an anxiety disorder and that the risk factors for mental problems among women reported to be related to hormonal and biochemical differences between both genders [3].

Ethnicity in the present study was significantly associated with depression and stress, with higher depression and stress levels reported in the minority ethnic groups followed by the Arab ethnic group, respectively. Higher depression and stress levels in the minority group could be attributed to the lack of expatriate communities from the same country or culture as the student himself. The detachment from their cultures with the lack of alternative community and support system deepens the feeling of loneliness and isolation from their surroundings. It was reported that minority ethnic groups may predispose to such psychological problems as minority students experience feeling of loneliness and discrimination [21]. While the high prevalence of depression and stress among the Arabs ethnic group may be related to the deteriorated states that their countries are having currently. The effects of such deterioration extend to affect those students' careers and personal lives.

There was no significant association between marital status and PD in the present study. However, it was reported in earlier studies that depression, anxiety, and stress were more likely to be higher among single students, and married students were less prone to PD [18], [19].

The present study demonstrated no significant association between depression, anxiety, and stress and monthly income of international students. This may be because this study was done on international students who are more inclined to manage their finances properly to avoid any unfavorable situations. In addition, reduced student prices are provided in many places in Malaysia including university canteens, public parks, tourist attractions and public transportation, etc. However, it was reported earlier that there was a significant association between socioeconomic level with stress and depression, as higher stress and depression scores were recorded in the low socioeconomic class [18], [19].

Study program was not associated with depression, anxiety, and stress among international students in the current study.

This might be since all international students are affected by the same circumstances regardless of the course of their studies. This is supported by an earlier study which found there were no significant associations between study program and depression, anxiety and stress scores [7]. However, other research findings reported that medical students showed higher prevalence and severity of stress and depression compared to their non-medical peers [19]. On the other hand, it was reported that the depression and anxiety levels were higher among students who were studying social and political sciences compared to those who were studying engineering, medical, or basic sciences [17].

TABLE VI
MULTIPLE REGRESSION MODEL FOR DEPRESSION, ANXIETY AND STRESS
WITH CHARACTERISTICS OF STUDY PARTICIPANTS

Variables	Depression $r^2 = 0.065$		Anxiety $r^2 = 0.052$		Stress $r^2 = 0.049$	
	Regression coefficient (b)	p-value	Regression coefficient (b)	p-value	Regression coefficient (b)	p-value
Age Group:						
18-20	-.023	.815	.749	.812	1.992	.519
21-24	.139	.047 S	4.625	.004 S	2.656	.086
25-35	Ref.		Ref.		Ref.	
Gender:						
Male	.274	.835	.789	.554	-.914	.484
Female						
Nationality:						
Arabs	6.471	.003 S	4.282	.052	6.279	.004 S
Asians	4.186	.079	3.143	.192	5.381	.023 S
Africans	Ref.		Ref.		Ref.	
Others	7.293	.048 S	2.653	.477	7.064	.054
Educational level:						
Undergraduate	.310	.896	.456	.849	1.861	.429
Postgraduate						
Study Program:						
Non-medical	.681	.670	-.605	.709	.037	.981
Medical/paramedical						
Monthly Income:						
< 2000 RM	-.029	.989	-1.559	.455	-.351	.863
2000 – 5000 RM	.204	.921	-1.435	.488	-1.052	.603
> 5000 RM	Ref.		Ref.		Ref.	
Marital Status:						
Married	-.637	.617	1.003	.436	1.145	.365
Single						

Although university students often demonstrate low levels of mental well-being, they show poor help-seeking behaviors for their mental health problems [9]. In this study, help-seeking behavior was found to be positively associated with depression, anxiety, and stress scores as students with higher PD reported higher help-seeking behavior from support services available at the university. While they also showed that students with higher mental health levels were more likely to seek help from informal sources such as friends and partners. Students with lower mental health levels were unlikely to seek help from both formal and informal sources. Moreover, the current study found that only one in 10 international students had sought psychological help from the support services available at the university. Professional help-seeking behavior among students was demonstrated to range from 15% to one third of students [22], [23]. However, it was reported that only 3.77% of students sought help from the

available university support services [9]. Even though university counselling services are free, most students would seek help from external sources [24]. This may be attributed to lack of awareness of the symptoms of their mental problems or the available mental support services. Also, mental health illness may be stigmatized in some cultures, and therefore, it is important to normalize international students' views towards mental health issues.

The present study found that 78.6% of international students were unaware of the presence of on-campus mental support services. This is consistent with previous research that found that international students underutilize mental health services, with 50% of international students being unaware of the presence of such services [25].

The limitations of the current study include that the design of the study was cross-sectional which cannot determine the incidence rate of PD - it can only determine its prevalence. Also, a non-probability sampling method was used, and the samples were taken without regard to the correct distribution characteristics of the international students at the university due to lack of epidemiological data for this population. This may introduce sampling bias and affect the generalizability of the study findings. Moreover, it is important to mention that the DASS-21 questionnaire can be used to assess prevalence and severity of PD and not to diagnose psychological disorders.

IV. CONCLUSION

In conclusion, this study revealed high prevalence and severity of PD among international students that were associated with age and ethnicity. Also, very poor mental help-seeking behavior and awareness of the available on-campus mental support services were reported. These findings highlight the importance of screening and taking proper interventions to improve the mental well-being of this vulnerable population.

REFERENCES

- [1] X. Lei, L. Xiao, Y. Liu, and Y. Li, "Prevalence of Depression among Chinese University Students: A Meta-Analysis," pp. 1–14, 2016.
- [2] Í. J. S. Ribeiro, R. Pereira, I. v Freire, B. G. de Oliveira, C. A. Casotti, and E. N. Boery, "Stress and Quality of Life Among University Students: A Systematic Literature Review," *Health Professions Education*, vol. 4, no. 2, pp. 70–77, 2018.
- [3] K. Macauley, L. Plummer, C. Bemis, G. Brock, C. Larson, and J. Spangler, "Prevalence and Predictors of Anxiety in Healthcare Professions Students," *Health Professions Education*, vol. 4, no. 3, pp. 176–185, 2018.
- [4] S. H. Ridner, "Psychological distress: concept analysis," 2004.
- [5] A. Drapeau, A. Marchand, and D. Beaulieu-prévost, "Epidemiology of Psychological Distress," 2007.
- [6] T. A. Rn and B. M. Gp, "Understanding persons with psychological distress in primary health care," pp. 19–21, 2016.
- [7] K. Shamsuddin et al., "Correlates of depression, anxiety and stress among Malaysian university students," *Asian Journal of Psychiatry*, vol. 6, no. 4, pp. 318–323, 2013.
- [8] J. A. Gold, B. Johnson, G. Leydon, R. M. Rohrbaugh, and K. M. Wilkins, "Mental Health Self-Care in Medical Students: A Comprehensive Look at Help-Seeking," pp. 37–46, 2015.
- [9] J. Goodwin, L. Behan, P. Kelly, K. Mccarthy, and A. Horgan, "Help-seeking behaviors and mental well-being of first year undergraduate university students," *Psychiatry Research*, vol. 246, pp. 129–135, 2016.

- [10] D. Sarokhani, A. Delpisheh, Y. Veisani, M. T. Sarokhani, R. E. Manesh, and K. Sayehmiri, "Prevalence of depression among university students: A systematic review and meta-analysis study," *Depression Research and Treatment*, vol. 2013, 2013.
- [11] F. Seyfi, K. C. Poudel, J. Yasuoka, K. Otsuka, and M. Jimba, "Intention to seek professional psychological help among college students in Turkey: influence of help-seeking attitudes," 2013.
- [12] S. Mori, "Addressing the Mental Health Concerns of International Students," vol. 78, pp. 17–24, 2000.
- [13] L. Bradley, G. Parr, W. Y. Lan, R. Bingi, and L. J. Gould, "Counselling expectations of international students," *International Journal for the Advancement of Counselling*, vol. 18, no. 1, pp. 21–31, 1995.
- [14] The International mobility of students in Asia and the Pacific. (2013). Retrieved from <https://unesdo>
- [15] P. F. Lovibond, "Pergamon The Structure of Negative Emotional States: Scales (DASS) with the Beck Depression and Anxiety inventories" vol. 33, no. 3, pp. 335–343, 1995.
- [16] J. D. Henry and J. R. Crawford, "The short-form version of the Depression anxiety stress scales (DASS-21): Construct validity and normative data in a large non-clinical sample," *British Journal of Clinical Psychology*, vol. 44, no. 2, pp. 227–239, 2005.
- [17] N. Bayram, N., Bilgel, "The prevalence and socio-demographic correlations of depression, anxiety and stress among a group of university students," pp. 667–672, 2008.
- [18] W. Yousif, A. Wahed, and S. K. Hassan, "Alexandria University Faculty of Medicine Prevalence and associated factors of stress , anxiety and depression among medical Fayoum University students," *Alexandria Journal of Medicine*, vol. 53, no. 1, pp. 77–84, 2017.
- [19] C. K. Teh, C. W. Ngo, R. Aniyah, R. Vellasamy, and K. Suresh, "Depression, Anxiety and Stress among Undergraduate Students: A Cross Sectional Study," no. November, pp. 260–268, 2015.
- [20] L. N. Dyrbye, M. R. Thomas, and T. D. Shanafelt, "Systematic Review of Depression, Anxiety, and Other Indicators of Psychological Distress among U. S. and Canadian Medical Students," vol. 81, no. 4, pp. 354–373, 2006.
- [21] L. Dyrbye, A. Eacker, and D. Power, "Race, Ethnicity, and Medical Student Well-being in the United States," no. April 2014, 2007.
- [22] F. I. Bilican, "Help-Seeking Attitudes and Behaviors Regarding Mental Health Among Turkish College Students Help-Seeking Attitudes and Behaviors Regarding Mental Health Among Turkish College Students," vol. 7411, 2014.
- [23] S. Quigley, E. Bainbridge, C. McDonald, and J. Lally, "Stigma of mental illness and help-seeking intention in university students," pp. 253–260, 2013.
- [24] N. J. Reavley, S. Cvetkovski, A. F. Jorm, and D. I. Lubman, "Help-seeking for substance use , anxiety and affective disorders among young people : results from the 2007 Australian National Survey of Mental Health and Wellbeing," 2010.
- [25] J. Lau, T. Garza, and H. Garcia, "International Students in Community Colleges : On-Campus Services Used International Students in Community Colleges : On- Campus Services Used and Its Affect on Sense of Belonging," *Community College Journal of Research and Practice*, vol. 00, no. 00, pp. 1–13, 2018.